

## AN EIGENVALUE PROBLEM FOR A NON-BOUNDED QUASILINEAR OPERATOR

JOSÉ CARMONA<sup>1</sup> AND ANTONIO SUÁREZ<sup>2</sup>

<sup>1</sup>*Departamento de Algebra y Análisis Matemático, Facultad de Ciencias,  
Cañada de San Urbano, Almería, Spain (jcarmona@ual.es)*

<sup>2</sup>*Departamento de Ecuaciones Diferenciales y Análisis Numérico,  
Facultad de Matemáticas, Sevilla, Spain (suarez@us.es)*

(Received 29 April 2003)

*Abstract* In this paper we study the eigenvalues associated with a positive eigenfunction of a quasilinear elliptic problem with an operator that is not necessarily bounded. For that, we use the bifurcation theory and obtain the existence of positive solutions for a range of values of the bifurcation parameter.

*Keywords:* bifurcation; positive solutions; non-bounded quasilinear operator; eigenvalue problems

2000 *Mathematics subject classification:* Primary 35J60; 35J25  
Secondary 35D05

### 1. Introduction

Let  $\Omega$  be a bounded open subset of  $\mathbb{R}^N$  with sufficiently smooth boundary  $\partial\Omega$  and let  $A(x, s)$  be a real symmetric matrix whose entries,  $a_{ij} : \bar{\Omega} \times \mathbb{R}_0^+ \rightarrow \mathbb{R}$ , are Carathéodory functions.

We assume that there exists a positive constant  $\alpha$  satisfying, for every  $(x, s, \xi) \in \Omega \times \mathbb{R}^+ \times \mathbb{R}^N$ ,

$$A(x, s)\xi \cdot \xi \geq \alpha|\xi|^2. \quad (A_1)$$

In this paper we analyse the nonlinear eigenvalue problem

$$\left. \begin{aligned} -\operatorname{div}(A(x, u)\nabla u) &= \lambda u, & x \in \Omega, \\ u &= 0, & x \in \partial\Omega, \end{aligned} \right\} \quad (P_\lambda)$$

where we say that  $\lambda$  is an eigenvalue for this problem if  $(P_\lambda)$  admits a positive and non-trivial solution, that is, if there exists  $u \in H_0^1(\Omega)$ ,  $u \geq 0$ ,  $u \not\equiv 0$ , such that  $A(x, u)\nabla u \in (L^2(\Omega))^N$  and

$$\int_{\Omega} A(x, u)\nabla u \cdot \nabla v = \lambda \int_{\Omega} uv, \quad \forall v \in H_0^1(\Omega).$$

In addition to interest itself in the study of  $(P_\lambda)$ , this kind of equation has been used to model a species inhabiting  $\Omega$  where its diffusion depends on the density of the species, which arises in more realistic models (see [3, 4] and references therein).

Problem  $(P_\lambda)$  is well known when  $A$  does not depend on  $s$ , i.e. when  $A(x, s) = B(x)$  with  $B = (b_{ij})$  and  $b_{ij} \in L^\infty(\Omega)$ ,  $b_{ij} \geq b_0 > 0$  in  $\Omega$ . In this case, there exists the principal eigenvalue, denoted by  $\lambda_1(B)$ , for the problem

$$\left. \begin{aligned} -\operatorname{div}(B(x)\nabla u) &= \lambda u, & x \in \Omega, \\ u &= 0, & x \in \partial\Omega, \end{aligned} \right\} \quad (1.1)$$

being the unique eigenvalue with a positive eigenfunction (see, for example, [6]).

In [2], assuming that  $A$  satisfies  $(A_1)$  and

$$|A(x, s)| \leq \beta \quad \text{for each } (x, s) \in \Omega \times \mathbb{R}, \quad (A_2)$$

the author proved that for each  $r > 0$ , there exists  $\lambda_r > 0$  and a positive solution  $u_r \in H_0^1(\Omega)$ , of  $(P_{\lambda_r})$  such that  $\|u_r\|_2 = r$ . Moreover, defining

$$\lambda_0 := \lambda_1(A(x, 0)),$$

he showed that if  $r \rightarrow 0$ , then  $\lambda_r \rightarrow \lambda_0$  and  $u_r/r$  converges to a positive eigenfunction associated with  $\lambda_0$  in  $H_0^1(\Omega)$ . Finally, if  $A$  also verifies

$$\lim_{s \rightarrow \infty} A(x, s) = A_\infty(x) \quad \text{uniformly in } x \in \Omega, \quad (A_3)$$

then  $\lambda_r \rightarrow \lambda_\infty$  and  $u_r/r$  goes to a positive eigenfunction associated with  $\lambda_\infty$  in  $H_0^1(\Omega)$  as  $r \rightarrow \infty$ , where

$$\lambda_\infty := \lambda_1(A_\infty(x)).$$

In [5], a slight modification of  $(P_\lambda)$  is analysed. Under conditions  $(A_1)$ – $(A_3)$ ,  $\lambda u + h(x)$  for some  $0 \leq h \in L^2(\Omega)$  is considered instead of  $\lambda u$ . But the arguments used to prove the existence of a solution leads to the trivial one in the case  $h \equiv 0$ .

In [1], assuming in addition the existence of an Osgood function  $\omega : \mathbb{R}_0^+ \rightarrow \mathbb{R}$  such that

$$|A(x, s_1) - A(x, s_2)| \leq \omega(|s_1 - s_2|), \quad (A_4)$$

for every  $(x, s_1), (x, s_2) \in \Omega \times \mathbb{R}$ , using a bifurcation analysis, the authors study a more general problem

$$\begin{aligned} -\operatorname{div}(A(x, u)\nabla u) &= f(\lambda, x, s), & x \in \Omega, \\ u &= 0, & x \in \partial\Omega, \end{aligned}$$

for  $f : \mathbb{R} \times \bar{\Omega} \times \mathbb{R} \mapsto \mathbb{R}$  and  $A$  satisfying  $(A_1)$ – $(A_4)$ . In the particular case  $f(\lambda, x, s) = \lambda s$ , from their results can be deduced the existence of an unbounded continuum (closed and connected subset) of positive solutions bifurcating from the trivial solution at  $\lambda = \lambda_0$  and meeting with infinity at the value  $\lambda = \lambda_\infty$ . Thus, as a consequence, there exists a positive

solution of  $(P_\lambda)$  for  $\lambda \in (\lambda_0, \lambda_\infty)$  or  $(\lambda_\infty, \lambda_0)$ . In the following section we complete this study for  $A$  satisfying  $(A_1)$ – $(A_4)$  by giving sufficient conditions for the uniqueness of a positive solution.

The main goal of this work (see §3) is to analyse  $(P_\lambda)$  when  $A$  is not necessarily bounded and/or does not satisfy  $(A_3)$ . In this case, we show that there exists an unbounded continuum of positive solutions bifurcating from the trivial one at  $\lambda = \lambda_0$ . If, in addition, there exists a continuous function  $g : \mathbb{R}_0^+ \rightarrow \mathbb{R}$ , with  $\lim_{s \rightarrow +\infty} g(s) = +\infty$ , satisfying, for every  $(x, s, \xi) \in \Omega \times \mathbb{R}^+ \times \mathbb{R}^N$ ,

$$A(x, s)\xi \cdot \xi \geq g(s)|\xi|^2 \geq \alpha|\xi|^2, \tag{A_\infty}$$

then the bifurcation from infinity at  $\lambda = \lambda_\infty$  (which exists in the bounded case) ‘disappears’. Specifically, there exists at least one positive solution  $u_\lambda$  for  $\lambda \in (\lambda_0, \infty)$  and  $\|u_\lambda\| \rightarrow \infty$  as  $\lambda \rightarrow \infty$ . However, if  $A$  is bounded in a subset of  $\Omega$ , then again a bifurcation to infinity exists.

Throughout the paper we will use the following notation.

(i)  $H_0^1(\Omega)$  and  $E = C_0(\bar{\Omega})$  are the usual Sobolev space and the space of the continuous functions in  $\bar{\Omega}$  vanishing on  $\partial\Omega$  endowed with the norms  $\|u\| = \|\nabla u\|_2$  and  $\|u\|_0 = \sup_\Omega |u|$ , respectively.

(ii)  $\text{cl}(D)$  denotes the closure of the set  $D$ .

(iii)  $\mathcal{S}$  denotes the set

$$\mathcal{S} = \text{cl}\{(\lambda, u) \in \mathbb{R} \times E : u \text{ is a solution for } (P_\lambda), u \geq 0, u \not\equiv 0\}.$$

Any continuum subset of  $\mathcal{S}$  will be called a continuum of positive solutions of  $(P_\lambda)$ , although it may contain the trivial solution  $(\lambda, 0)$  for some value of  $\lambda > 0$ .

(iv)  $I$  will denote both the identity matrix and the identity operator.

(v) Given square matrices  $B_1, B_2$  we say that  $B_1 > 0$  (respectively,  $B_1 \geq 0$ ) if the quadratic form induced by  $B_1$  is definite positive (respectively, semidefinite positive). We say that  $B_1 < B_2$  (respectively,  $B_1 \leq B_2$ ) if  $B_2 - B_1 > 0$  (respectively,  $B_2 - B_1 \geq 0$ ).

(vi) The map  $\text{Proj}_\mathbb{R} : \mathbb{R} \times E \mapsto \mathbb{R}$  stands for the projection of the product space  $\mathbb{R} \times E$  onto  $\mathbb{R}$ .

**2. The case of bounded matrices A**

In order to study problem  $(P_\lambda)$ , let us recall that, for matrices  $A$  satisfying  $(A_1), (A_2)$ , if  $u \in H_0^1(\Omega)$  is a solution of  $(P_\lambda)$ , then using the De Giorgi–Stampacchia Theorem (see [9, Théorème 7.3] and [7, Theorem I] or [8, Theorem 8.29]),  $u \in C^{0,\gamma}(\bar{\Omega})$  for some  $0 < \gamma < 1$ . Moreover, if the coefficients of the matrix  $A$  satisfy

$$a_{ij} \in C^{1,\gamma'}(\bar{\Omega} \times \mathbb{R}) \quad \text{for some } 0 < \gamma' < 1, \tag{2.1}$$

then by Theorem 15.17 in [8] we have that  $u \in C_0^{2,\gamma\gamma'}(\bar{\Omega})$ .

We also recall that for every  $(\lambda, u) \in \mathcal{S}$  with  $u \in C^1(\bar{\Omega})$  and  $u \not\equiv 0$ , using the Hopf maximum principle, we have that  $u > 0$  in  $\Omega$  and the normal exterior derivative  $\partial u / \partial n_e$  is negative in  $\partial\Omega$ .

The following lemma gives us necessary conditions in  $\lambda \in \mathbb{R}$  for which  $(P_\lambda)$  admits a solution in some special cases.

**Lemma 2.1.** *Assume  $(A_1)$ ,  $(A_3)$  and that  $(P_\lambda)$  admits a positive solution. Then*

- (1)  $\lambda_0 \leq \lambda$  (respectively,  $<$ ,  $\geq$ ,  $>$ ) if for every  $s \in \mathbb{R}^+$ ,  $A(x, 0) \leq A(x, s)$  (respectively,  $<$ ,  $\geq$ ,  $>$ ); and
- (2)  $\lambda_\infty \geq \lambda$  (respectively,  $>$ ,  $\leq$ ,  $<$ ) if for every  $s \in \mathbb{R}^+$ ,  $A_\infty(x) \geq A(x, s)$  (respectively,  $>$ ,  $\leq$ ,  $<$ ).

**Proof.** The result follows from the fact that for given symmetric matrices  $B_1(x)$ ,  $B_2(x)$  for which there exist  $\lambda_1(B_1)$  and  $\lambda_1(B_2)$ , with  $0 < B_1 \leq B_2$ , then

$$\lambda_1(B_1) = \inf \left\{ \int_{\Omega} B_1(x) \nabla u \cdot \nabla u, u \in H_0^1(\Omega), \|u\|_2 = 1 \right\} \leq \lambda_1(B_2).$$

Thus, if  $u \in H_0^1(\Omega)$  is a solution of  $(P_\lambda)$ , we conclude by taking into account that  $\lambda = \lambda_1(A(x, u))$ .  $\square$

The main result of this section is the following.

**Theorem 2.2.** *Assume  $(A_1)$ – $(A_4)$ . We have that  $\lambda_0$  and  $\lambda_\infty$  are the only bifurcation points from the trivial solution and from infinity, respectively, and there exists a continuum  $\Sigma \subset \mathcal{S}$  of positive solutions meeting  $(\lambda_0, 0)$  and  $(\lambda_\infty, \infty)$ ; in particular,  $(P_\lambda)$  possesses a positive solution for every  $\lambda \in (\lambda_0, \lambda_\infty)$  or  $\lambda \in (\lambda_\infty, \lambda_0)$ . Moreover,*

- (i) *the bifurcation from  $\lambda_0$  is subcritical (respectively, supercritical) if there exists  $s_0 > 0$  such that*

$$A(x, s) < A(x, 0) \quad (\text{respectively, } A(x, s) > A(x, 0)), \quad \forall s \in (0, s_0),$$

- (ii) *the bifurcation from  $\lambda_\infty$  is subcritical (respectively, supercritical) if*

$$A(x, s) < A_\infty(x) \quad (\text{respectively, } A(x, s) > A_\infty(x)), \quad \forall s \in \mathbb{R}^+.$$

Furthermore, we have the following.

- (i) *If  $A(x, 0) < A(x, s) < A_\infty(x)$  for every  $s \in \mathbb{R}^+$ , then there exists a non-trivial solution for  $(P_\lambda)$  if, and only if,  $\lambda \in (\lambda_0, \lambda_\infty)$ ; in particular,  $\text{Proj}_{\mathbb{R}} \Sigma = [\lambda_0, \lambda_\infty)$ . If, in addition,  $A(x, s)$  is increasing in  $s$  and it satisfies (2.1), the solution is unique.*
- (ii) *If  $A(x, 0) > A(x, s) > A_\infty(x)$  for every  $s \in \mathbb{R}^+$ , then there exists a non-trivial solution for  $(P_\lambda)$  if, and only if,  $\lambda \in (\lambda_\infty, \lambda_0)$ ; in particular,  $\text{Proj}_{\mathbb{R}} \Sigma = (\lambda_\infty, \lambda_0]$ .*

**Proof.** The existence of the continuum  $\Sigma$  of positive solutions follows by Theorem 5.1 in [1]; in particular, we have the existence of positive solutions for every  $\lambda$  in  $(\lambda_0, \lambda_\infty)$  or in  $(\lambda_\infty, \lambda_0)$ .

The description  $\text{Proj}_{\mathbb{R}} \Sigma$ , in the cases  $A(x, 0) < A(x, s) < A_\infty(x)$  or  $A(x, 0) < A(x, s) < A_\infty(x)$  for every  $s \in \mathbb{R}^+$ , follows directly from Lemma 2.1. Moreover, arguing as in that lemma we get the laterality of the bifurcations.

Now, assume that  $A(x, s)$  is increasing in  $s$  and (2.1) is satisfied. In order to prove the uniqueness of the solution for  $(P_\lambda)$ , let us suppose that there exists  $\lambda \in (\lambda_0, \lambda_\infty)$  for which  $(P_\lambda)$  admits two solutions,  $u_1, u_2 \in E$ , with  $u_1 \not\equiv u_2$ . We claim that  $u_1, u_2$  can be chosen such that  $u_1 \leq u_2$ . Indeed, this is a consequence of the existence of a sequence  $(\lambda_n, u_n)$  with  $\lambda_n \rightarrow \lambda_0$  and  $u_n \rightarrow 0$  in  $E$ . In fact, by regularity results,  $u_n \rightarrow 0$  in  $C^1(\Omega)$ . Thus, for  $\lambda_n < \lambda$ ,  $u_n$  is a subsolution for  $(P_\lambda)$  and for large  $n$ ,  $u_n \leq \min\{u_1, u_2\}$ . Then, by the subsolution and supersolution method, there exists  $w \in E$ , a solution of  $(P_\lambda)$  with

$$u_n \leq w \leq u_1, \quad u_n \leq w \leq u_2.$$

This implies that  $w \not\equiv u_1$  or  $w \not\equiv u_2$ , and the claim is proved by taking  $u_1 = w$  and  $u_2 = u_i$  for some  $i = 1, 2$ .

Now we take  $v = u_2^2/u_1$  as a test function in the equation satisfied by  $u_1$ , and  $v = u_2$  in that satisfied by  $u_2$ . Thus, subtracting both equalities we have

$$\begin{aligned} 0 &= \int_{\Omega} A(x, u_1) \nabla u_1 \cdot \nabla \left( \frac{u_2^2}{u_1} \right) - \int_{\Omega} A(x, u_2) \nabla u_2 \cdot \nabla u_2 \\ &= - \int_{\Omega} A(x, u_1) \left( \frac{u_2}{u_1} \nabla u_1 - \nabla u_2 \right) \cdot \left( \frac{u_2}{u_1} \nabla u_1 - \nabla u_2 \right) \\ &\quad - \int_{\Omega} (A(x, u_2) - A(x, u_1)) \nabla u_2 \cdot \nabla u_2 < 0. \end{aligned}$$

This contradiction gives the uniqueness. □

### 3. The case of unbounded matrices A

In this section, we study  $(P_\lambda)$  when  $A$  is not necessarily bounded and does not satisfy  $(A_3)$ . We prove firstly that every solution of  $(P_\lambda)$  is bounded. More precisely we have the following lemma.

**Lemma 3.1.** *Let  $A(x, s)$  satisfy  $(A_1)$  and let  $u \in H_0^1(\Omega)$  be a solution of  $(P_\lambda)$ , then  $u \in E$ . Moreover, there exist positive constants  $c_1, c_2, \gamma_1, \gamma_2$  such that*

$$\|u\|_0^{\gamma_1} \leq c_1 + c_2 \|u\|^{\gamma_2}. \tag{3.1}$$

**Proof.** Once we know that  $u \in L^\infty(\Omega)$ , and  $\|u\|_\infty^{\gamma_1} \leq c_1 + c_2 \|u\|^{\gamma_2}$  for some positive constants  $c_1, c_2, \gamma_1, \gamma_2$ , then the result follows directly from the De Giorgi–Stampacchia Theorem. Let us prove the  $L^\infty(\Omega)$ -estimate. We consider for every  $k \in \mathbb{R}^+$  the function  $G_k : \mathbb{R}_0^+ \rightarrow \mathbb{R}_0^+$  given by

$$G_k(s) = \begin{cases} 0, & 0 \leq s \leq k, \\ s - k, & s > k. \end{cases}$$

Thus, we can take  $v = G_k(u)$  as a test function in the weak equation satisfied by  $u$  and using  $(A_1)$  we have

$$\alpha \|\nabla G_k(u)\|_2^2 \leq \int_{\Omega} A(x, u) \nabla u \nabla G_k(u) \leq \lambda \int_{\Omega_k} u G_k(u), \tag{3.2}$$

where  $\Omega_k \equiv \{x \in \Omega : u(x) > k\}$ .

Using the Sobolev and Hölder inequalities, in the case  $N > 2$ , by (3.2) we have, for  $u \in L^r(\Omega)$  with  $r > 2^*/(2^* - 1)$  and for some positive constant  $c$ ,

$$\|G_k(u)\|_{2^*}^2 \leq c \|u\|_r \|G_k(u)\|_{2^*} (\text{meas } \Omega_k)^{(1-1/r-1/2^*)}. \tag{3.3}$$

Taking into account that, for every  $h > k$ ,  $G_k(u) \geq h - k$  in  $\Omega_h$ , (3.3) implies that

$$(h - k)(\text{meas } \Omega_h)^{1/2^*} \leq c \|u\|_r (\text{meas } \Omega_k)^{(1-1/r-1/2^*)},$$

or, equivalently,

$$\text{meas } \Omega_h \leq \frac{c \|u\|_r^{2^*} (\text{meas } \Omega_k)^{2^*-1-2^*/r}}{(h - k)^{2^*}}. \tag{3.4}$$

We can now apply the Stampacchia Lemma [9, Lemma 4.1] to deduce that

- (i) if  $u \in L^r(\Omega)$  with  $r > N/2$ , then  $u \in L^\infty(\Omega)$  and  $\|u\|_\infty \leq c \|u\|_r$ ;
- (ii) if  $u \in L^r(\Omega)$  with  $r = N/2$ , then  $u \in L^t(\Omega)$  for  $t \in [1, \infty)$  and  $\|u\|_t^t \leq c + c' \|u\|_r^t$ ;  
and
- (iii) if  $u \in L^r(\Omega)$  with  $r < N/2$ , then  $u \in L^t(\Omega)$  for

$$t = \frac{2^* r}{(2 - 2^*)r + 2^*} - \delta$$

and  $\delta > 0$  arbitrarily small—moreover,  $\|u\|_t^t \leq c + c' \|u\|_r^{t+\delta}$ .

Since  $u \in L^{2^*}(\Omega)$  and  $2^* > 2^*/(2^* - 1)$ , we can argue as before for  $r_0 = 2^*$ . Thus, if  $2^* > N/2$ , we obtain the  $L^\infty(\Omega)$  estimate by using item (i) above. In the case  $2^* = N/2$  we use item (ii) in order to take  $r_1 > N/2$  and again the  $L^\infty(\Omega)$  estimate follows from item (i) above. Finally, in the case  $2^* < N/2$  we can take

$$r_1 = \frac{2^* r_0}{(2 - 2^*)r_0 + 2^*} - \delta_1 > r_0.$$

As before, if  $r_1 \geq N/2$ , we easily conclude. In the other case we take

$$r_2 = \frac{2^* r_1}{(2 - 2^*)r_1 + 2^*} - \delta_2.$$

By an iterative argument we conclude after a finite number of steps. Indeed, in the other case, we have that  $r_n$  is bounded, where  $r_n$  is defined recurrently by

$$r_0 = 2^*,$$

$$r_{n+1} = \frac{2^* r_n}{(2 - 2^*)r_n + 2^*} - \delta_{n+1},$$

where  $\lim_{n \rightarrow \infty} \delta_n = 0$ . Moreover,  $r_n$  is non-decreasing and so it converges to  $r \in (2^*, N/2]$ , which satisfies

$$r = \frac{2^*r}{(2 - 2^*)r + 2^*};$$

that is,  $2^* = (2 - 2^*)r + 2^*$ , which implies that  $r = 0$ , and this is a contradiction.

Observe that the estimate (3.1) follows, after this finite number of steps, from estimates in items (i)–(iii) and the Sobolev embedding.

Finally, in the case  $N = 2$  we can choose  $r > q/(q - 2)$  for any  $q > 2$  and argue as before with  $2^*$  replaced by  $q$ . In this case we obtain the  $L^\infty(\Omega)$  estimate directly by using item (i) above.  $\square$

In this section we assume, instead of  $(A_2)$ , that for each  $s_0 \in \mathbb{R}^+$  there exists  $\beta(s_0)$  such that

$$|A(x, s)| \leq \beta(s_0), \tag{\tilde{A}_2}$$

for  $(x, s) \in \bar{\Omega} \times [0, s_0]$ .

We consider the truncated problems

$$\left. \begin{aligned} -\operatorname{div}(A(x, T_n(u))\nabla u) &= \lambda u, & x \in \Omega, \\ u &= 0, & x \in \partial\Omega, \end{aligned} \right\} \tag{P_{\lambda,n}}$$

$T_n(s)$  being the map defined, for each  $n \in \mathbb{N}$ , by

$$T_n(s) = \begin{cases} s, & 0 \leq s \leq n, \\ n, & s > n. \end{cases}$$

By Theorem 2.2, there exist  $\Sigma_n$  unbounded maximal continua of positive solutions such that  $(\lambda_0, 0) \in \Sigma_n$  for each  $n \in \mathbb{N}$ . Now, we can prove the following theorem.

**Theorem 3.2.** *Suppose that  $A$  satisfies  $(A_1)$ ,  $(A_4)$  and  $(\tilde{A}_2)$ . Then there exists an unbounded continuum  $\Sigma \subset \mathcal{S}$  such that  $(\lambda_0, 0) \in \Sigma$ .*

**Proof.** Firstly, we denote by  $\Sigma_k^n$  the connected component of  $\Sigma_k \cap (\mathbb{R} \times \bar{B}_n(0))$  containing  $(\lambda_0, 0)$ . We claim that

$$\Sigma_k^n = \Sigma_n^n \quad \text{for } k \geq n. \tag{3.5}$$

Indeed, if  $k \geq n$  and  $(\lambda, u) \in \Sigma_k^n$ , then  $u$  is a solution of  $(P_{\lambda,n})$ . Thus,  $\Sigma_k^n$  is a closed and connected subset of

$$\operatorname{cl}\{(\lambda, u) \in \mathbb{R} \times E : u \text{ is a non-trivial solution of } (P_{\lambda,n})\}$$

containing  $(\lambda_0, 0)$ . So,  $\Sigma_k^n \subset \Sigma_n^n$ , from which we deduce that  $\Sigma_k^n \subset \Sigma_n^n$ . We can reason similarly and obtain that  $\Sigma_n^n \subset \Sigma_k \cap (\mathbb{R} \times \bar{B}_n(0))$ ; thus  $\Sigma_n^n$  and  $\Sigma_k^n$  are connected components of  $\Sigma_k \cap (\mathbb{R} \times \bar{B}_n(0))$  containing  $(\lambda_0, 0)$ , which implies (3.5). So, we get

$$\Sigma_n^n = \lim_k \Sigma_k^n.$$

Therefore, for each  $n \in \mathbb{N}$  we have a continuum

$$\Sigma_n^n \subset \text{cl}\{(\lambda, u) \in \mathbb{R} \times E : u \text{ is a non-trivial solution of } (P_\lambda)\}$$

containing  $(\lambda_0, 0)$ , and if  $(\lambda, u) \in \Sigma_n^n$ , then  $\|u\|_0 \leq n$ .

Now, we are going to prove that

$$\Sigma_n^n \subset \Sigma_{n+1}^{n+1} \quad \text{for each } n \in \mathbb{N}. \quad (3.6)$$

Indeed, observe that

$$\Sigma_n^n = \Sigma_{n+1}^n \subset \Sigma_{n+1} \cap (\mathbb{R} \times \bar{B}_n(0)) \subset \Sigma_{n+1} \cap (\mathbb{R} \times \bar{B}_{n+1}(0)),$$

so, since  $\Sigma_{n+1}^{n+1}$  is the connected component of  $\Sigma_{n+1} \cap (\mathbb{R} \times \bar{B}_{n+1}(0))$  containing  $(\lambda_0, 0)$  and  $\Sigma_n^n$  is a connected component of such a subset containing it, (3.6) follows.

Finally, we show that the set

$$\Sigma = \bigcup_{n=1}^{\infty} \Sigma_n^n$$

satisfies the theorem. Firstly, observe that since  $\Sigma_n^n$  is unbounded,  $\Sigma$  is also unbounded. Indeed, since  $\text{Proj}_{\mathbb{R}} \Sigma_n^n$  is bounded, so there exists a connected subset of  $\Sigma_n^n \cap (\mathbb{R} \times \bar{B}_n(0))$  containing  $(\lambda_0, 0)$  and intersecting with  $\mathbb{R} \times \partial \bar{B}_n(0)$  for each  $n \in \mathbb{N}$ ; i.e. for each  $n \in \mathbb{N}$  there exists  $(\lambda_n, u_n) \in \Sigma_n^n$ , with  $\|u_n\|_0 = n$ .

On the other hand, since  $\Sigma_n^n$  is connected and  $(\lambda_0, 0) \in \Sigma_n^n$  for each  $n \in \mathbb{N}$ , it follows that  $\Sigma$  is connected.

Finally, we will prove that  $\Sigma$  is closed. Let  $(\lambda, u) \in \bar{\Sigma}$ . Since  $\bar{\Sigma}$  is connected, there exists a connected and bounded set  $\Sigma' \subset \bar{\Sigma}$  containing  $(\lambda_0, 0)$  and  $(\lambda, u)$ . Thus, there exists  $n \in \mathbb{N}$  such that

$$\Sigma' \subset \text{cl}\{(\lambda, u) \in \mathbb{R} \times E : \|u\|_0 \leq n, u \text{ is a non-trivial solution of } (P_{\lambda, n})\}.$$

In particular,  $\Sigma' \subset \Sigma_n^n \cap (\mathbb{R} \times \bar{B}_n(0))$ , from which  $\Sigma' \subset \Sigma_n^n$  and so  $(\lambda, u) \in \Sigma_n^n \subset \Sigma$ .  $\square$

**Remark 3.3.**

- (i) We point out that the above result is true even in the case in which the limit of  $A(x, s)$  does not exist as  $s \rightarrow \infty$ .
- (ii) In the case where  $A$  is bounded in some subset of  $\Omega$ , we can conclude that  $\text{Proj}_{\mathbb{R}} \Sigma$  is bounded. Indeed, assume that  $|A(x, s)| \leq \gamma$  if  $x \in B$ , where  $B$  is a ball such that  $B \subset \Omega$ , then using the monotonicity of the principal eigenvalue with respect to the domain, we obtain

$$\lambda = \lambda_1(A(x, u)) \leq \lambda_1^B(A(x, u)) \leq \lambda_1^B(\gamma I) = \gamma \lambda_1^B(I).$$

- (iii) In this case we can obtain a similar result to the main one in [2]. Indeed, for each  $r > 0$  there exists  $\lambda_r > 0$  and  $u_r \in H_0^1(\Omega)$ , a solution of  $(P_{\lambda_r})$  with  $\|u_r\|_0 = r$ .



In the next result we show that when  $A(x, s)$  tends to infinity as  $s \rightarrow \infty$  in the sense of  $(A_\infty)$ , then the bifurcation at infinity disappears, in some sense  $\lambda_\infty \rightarrow +\infty$  when  $A(x, s)$  tends to infinity.

**Theorem 3.4.** *Assume that  $A$  satisfies  $(A_4)$ ,  $(\tilde{A}_2)$  and  $(A_\infty)$ . Then there exists a continuum  $\Sigma \subset \mathcal{S}$  such that  $(\lambda_0, 0) \in \Sigma$ . Moreover, the interval  $(\lambda_0, +\infty) \subset \text{Proj}_{\mathbb{R}} \Sigma$  and*

$$\lim_{\substack{\lambda \rightarrow +\infty \\ (\lambda, u_\lambda) \in \Sigma}} \|u_\lambda\|_0 = +\infty.$$

**Proof.** The existence of the continuum unbounded  $\Sigma$  bifurcating from  $(\lambda_0, 0)$  follows by Theorem 3.2. Since  $\lambda = \lambda_1(A(x, u)) \geq \lambda_1(\alpha I) = \alpha \lambda_1(I)$ , there do not exist positive solutions for  $\lambda$  small. So, it suffices to prove that bifurcation from infinity is not possible. In order to do this, we observe that problem  $(P_\lambda)$  can be written as

$$\left. \begin{aligned} -\operatorname{div}(B(x, u)g(u)\nabla u) &= \lambda u, & x \in \Omega, \\ u &= 0, & x \in \partial\Omega, \end{aligned} \right\} \tag{P_\lambda}$$

where  $g$  is given by hypothesis  $(A_\infty)$  and

$$B(x, u) := \frac{A(x, u)}{g(u)}.$$

Moreover, if we perform the change of variable

$$w = \tilde{g}(u) = \int_0^u g(t) dt,$$

problem  $(P_\lambda)$  is equivalent to

$$\left. \begin{aligned} -\operatorname{div}(C(x, w)\nabla w) &= \lambda f(w), & x \in \Omega, \\ w &= 0, & x \in \partial\Omega, \end{aligned} \right\} \tag{Q_\lambda}$$

where

$$C(x, w) := B(x, \tilde{g}^{-1}(w)) \quad \text{and} \quad f(w) := \tilde{g}^{-1}(w).$$

Now we argue by contradiction, and assume that there exists a sequence of solutions  $(\lambda_n, u_n)$  of  $(P_{\lambda_n})$  such that  $\lambda_n \rightarrow \bar{\lambda} > 0$  and  $\|u_n\|_0 \rightarrow \infty$ . Then, by (3.1), we have that  $\|u_n\| \rightarrow \infty$  and taking  $w_n = \tilde{g}(u_n)$ , it is clear that  $\|w_n\|_0 \rightarrow \infty$ . In addition, since  $(A_\infty)$  implies that  $\alpha^2 \|u_n\|^2 \leq \|w_n\|^2$ , we also have that  $\|w_n\| \rightarrow \infty$ . For the normalized sequence  $z_n := w_n/\|w_n\|$  we know the existence of  $z \in H_0^1(\Omega)$ , such that

$$z_n \rightarrow z \quad \text{strongly in } L^2(\Omega) \text{ and a.e. in } \Omega.$$

and so, taking  $w_n/\|w_n\|^2$  as a test function in  $(Q_{\lambda_n})$ , we obtain that

$$\alpha \leq \int_\Omega C(x, w_n)\nabla z_n \cdot \nabla z_n = \lambda_n \int_\Omega \frac{f(w_n)}{\|w_n\|} z_n. \tag{3.7}$$

Now, taking into account that

$$\frac{f(s)}{s} \rightarrow 0 \quad \text{as } s \rightarrow \infty$$

and that  $f(s) \leq (1/\alpha)s$  for each  $s \in \mathbb{R}^+$ , we can argue as in Theorem 5.5 in [1] and conclude that

$$\int_{\Omega} \frac{f(w_n)}{\|w_n\|} z_n \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Indeed, we can write for every  $n \in \mathbb{N}$

$$\begin{aligned} \int_{\Omega} \frac{f(w_n)}{\|w_n\|} z_n &= \int_{\Omega} \frac{f(w_n)}{\|w_n\|} (z_n - z) + \int_{\Omega} \frac{f(w_n)}{\|w_n\|} z \\ &\leq \frac{1}{\alpha} \|z_n\|_2 \|z_n - z\|_2 + \int_{\Omega_0} \frac{f(w_n)}{\|w_n\|} z, \end{aligned}$$

where  $\Omega_0 = \{x \in \Omega : z(x) \neq 0\}$ . Thus, we only have to prove that

$$\lim_{n \rightarrow \infty} \int_{\Omega_0} \frac{f(w_n)}{\|w_n\|} z = 0,$$

which is a direct consequence of the Lebesgue Theorem, since for a.e.  $x \in \Omega_0$ ,  $w_n(x) = z_n(x)\|w_n\| \rightarrow +\infty$  and then

$$\frac{f(w_n(x))}{\|w_n\|} z(x) \rightarrow 0 \quad \text{a.e. } x \in \Omega_0.$$

Thus, taking limits in (3.7), we have that  $\alpha \leq 0$ , which is a contradiction.  $\square$

**Acknowledgements.** This work was developed while J.C. was visiting the Departamento de Ecuaciones Diferenciales y Análisis Numérico, Universidad de Sevilla. He gratefully acknowledges the whole department for the warm hospitality and the friendly atmosphere. J.C. was partly supported by DGES Ministerio de Educación y Ciencia (Spain) no. PB98-1283. A.S. was supported by the Spanish Ministry of Science and Technology under grant BFM2000-0797.

## References

1. D. ARCOYA, J. CARMONA AND B. PELLACCI, Bifurcation for some quasilinear operators, *Proc. R. Soc. Edinb. A* **131** (2001), 733–765.
2. L. BOCCARDO, Positive eigenfunctions for a class of quasi-linear operator, *Boll. UMI B* **18** (1981), 951–959.
3. R. S. CANTRELL AND C. COSNER, Diffusive logistic equations with indefinite weights: population models in disrupted environments, Part I, *Proc. R. Soc. Edinb. A* **112** (1989), 293–318.
4. R. S. CANTRELL AND C. COSNER, Diffusive logistic equations with indefinite weights: population models in disrupted environments, Part II, *SIAM J. Math. Analysis* **22** (1991), 1043–1064.

5. J. CHABROWSKI, Some existence theorems for the Dirichlet problem for quasilinear elliptic equations, *Annli Mat. Pura Appl.* **158** (1991), 391–398.
6. D. G. DE FIGUEIREDO, *Positive solutions of semilinear elliptic problems*, Lectures Notes in Mathematics, vol. 957 (Springer, 1982).
7. E. DE GIORGI, *Sulla differenziabilità e l'analiticità degli estremali degli integrali multipli regolari*, Memorie della Accademia delle Scienze di Torino, pp. 25–43 (1957).
8. D. GILBARG AND N. TRUDINGER, *Elliptic partial differential equations of second order* (Springer, 1983).
9. G. STAMPACCHIA, Le problème de Dirichlet pour les équations elliptiques du second ordre à coefficients discontinus, *Annls Inst. Fourier* **117** (1965), 138–152.